

CONTAINS:

Cyromazine: 2.0% w/w (20 g/Kg)

Apply Hoko Ex at intervals exceeding 14 days. Animals may be present during applications. Often, 6-week intervals suffice.

Apply Hoko Ex at a dose of 250g/10m², independent of the method of application.

Apply Hoko Ex either as dry granules by dispersal if the manure is liquid or very moist, or dilute 250 g in 1-4 liter of water for spraying, or 10 liter water for pouring using a watering can. In case the manure is dry apply by pouring only.

The simultaneous use of an insecticide specific to adult flies, particularly coinciding with the first application of Hoko Ex, can significantly enhance fly control.

MANUFACTURER:

Hoko sarl.
15 Ave de Joinville
FRANCE – 94130 Nogent-sur-Marne
Tel: +41 31 302 8404
Fax: +41 31 302 8410
www.hoko.com

MARKETING AND SALES COMPANY:

Expert-Medical Ltd.
HUNGARY - 2072, Zsámbék, Majális köz 1.
Tel: +36 30 617 1692
expert@expert-medical.hu
www.expert-medical.hu

Quantity:
5 and 25 kg




HokoEx[®]

*Hoko Ex soluble granules
Insect Growth Regulator*

Hoko Ex is a larvicide used to control fly larvae in animal husbandry. Contains a potent growth disruptor specific to all fly species.

25 g / m²



In rearing facilities for **CATTLE** and **HORSES** apply Hoko Ex in a 50 cm band along the walls, under and next to machinery installations by pouring or spraying. When treating liquid manure the granules can be dispersed directly. The first application is best carried our 1-3 days after dung removal. In cow sheds with slatted floors apply Hoko Ex uniformly to the entire surface.

In **PIG** stys with completely slatted floors apply Hoko Ex to the entire surface, in the case of partially slatted floors apply to the slatted sections and the borders of the rest areas. Apply either by dispersal of dry granules, or pouring, or spraying. If possible treat before housing the pigs, shortly after manure or slurry removal is best.

In **POULTRY** houses with a slurry reservoir and in the presence of moist or liquid manure, disperse Hoko Ex granules directly, in dry manure situations use the pouring or spraying method. Apply to the entire surface. Commence treatment approximately 1 week after removal of excrement but before a depth of 10cm has accumulated.

Hoko Ex can also be used in sheep and goat sheds, rabbit farms, zoological gardens in the same manner.



OKO

www.hoko.hu

www.hoko.com • www.expert-medical.hu



Expert-Medical Ltd.
Always delivers the best solutions